

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/331107465>

An Innovative Home-Made Beebread Collector as a Tool for Sampling and Harvesting

Article in *Bee World* · January 2019

DOI: 10.1080/0005772X.2018.1556905

CITATIONS

3

READS

845

6 authors, including:



Giovanni Formato

Istituto Zooprofilattico Sperimentale del Lazio e Toscana

95 PUBLICATIONS 370 CITATIONS

[SEE PROFILE](#)



Marco Pietropaoli

L'Istituto Zooprofilattico Sperimentale del Lazio e Toscana

52 PUBLICATIONS 128 CITATIONS

[SEE PROFILE](#)



Norman Carreck

University of Sussex

175 PUBLICATIONS 4,033 CITATIONS

[SEE PROFILE](#)



Jozef Van der Steen

Alveus AB Consultancy

122 PUBLICATIONS 1,957 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



COLOSS monitoring of honey bee colony losses [View project](#)



Insignia [View project](#)

An Innovative Home-Made Beebread Collector as a Tool for Sampling and Harvesting

Giulio Loglio, Giovanni Formato, Marco Pietropaoli, Riccardo Jannoni-Sebastianini, Norman Carreck & Jozef van der Steen

To cite this article: Giulio Loglio, Giovanni Formato, Marco Pietropaoli, Riccardo Jannoni-Sebastianini, Norman Carreck & Jozef van der Steen (2019) An Innovative Home-Made Beebread Collector as a Tool for Sampling and Harvesting, *Bee World*, 96:1, 16-18, DOI: [10.1080/0005772X.2018.1556905](https://doi.org/10.1080/0005772X.2018.1556905)

To link to this article: <https://doi.org/10.1080/0005772X.2018.1556905>



Published online: 14 Feb 2019.



Submit your article to this journal [↗](#)



Article views: 11



View Crossmark data [↗](#)

An Innovative Home-Made Beebread Collector as a Tool for Sampling and Harvesting

Giulio Loglio, Giovanni Formato , Marco Pietropaoli , Riccardo Jannoni-Sebastianini, Norman Carreck and Jozef van der Steen



Introduction

A honey bee colony represents a practical tool for bio-indication, as it collects material from a large area while foraging for food (Grodzinski & Yorks, 1981; Porrini, Ghini, & Girotti, 2002; Raes, Cornelis, & Rzeznik, 1992; van der Steen, 2016; van der Steen, Cornelissen, Blacquiere, Pijnenburg, & Severijnen, 2016). There are several matrices available for sampling from the hive: nectar, honey, pollen, beebread, bees, and wax. Beebread as a food differs from corbiculate pollen loads in being fermented bee pollen (De Grandi-Hoffman, Chen, & Simonds, 2013; Fuenmayor et al., 2014) and it includes pollen, honey, and secretions of bees' salivary glands (Barajas, Cortes-Rodriguez, & Rodriguez-Sandoval, 2012; Vásquez & Olofsson, 2009) (Figure 1). In general, it has a higher nutritional value than pollen, better digestibility, a richer chemical composition (Anderson et al., 2014; Carroll et al., 2017; Habryka, Kruczek, & Drygaś, 2016) and is better absorbed thanks to the partial fermentation of components (Barene, Daberte, & Siksná, 2015).

Beebread contains 14–37% proteins, 24–34% carbohydrates, 3.2% lactic acid, 6–13% lipids, and 2.7% cellulose (Anđelković et al., 2012; Barene et al., 2015; Campos et al., 2008; De Grandi-Hoffman et al., 2013; Estevinho, Afonso, & Feás, 2011; Fuenmayor et al., 2014; Karmakar, 2015; Nagai, Nagashima, Myoda, & Inoue, 2004; Zuluaga, Serrato, & Quicazan, 2015). It further contains mineral components, vitamin K, and free amino acids (Gilliam, 1979; Nagai et al., 2004).

Mainly, technical reasons limit the use and commercialization of beebread. When beebread is intended to be used for research, medicinal, cosmetic or food purposes, it is necessary to collect it by hand, since this is the only way to guarantee the highest microbiological or chemical purity (Kieliszek et al., 2017). In this case, beebread is extracted manually, breaking the wax comb at room temperature or after freezing; using a large-mesh sieve, it is possible to collect the beebread in the form of small cylinders, separating them from the wax. It is possible to speed up the harvesting process using a strong flow of air that separates wax from the beebread. Otherwise, operators may use small metal or plastic tools to extract the beebread from the combs and send it to a laboratory. This procedure crumbles beebread and may not provide a clean sample, as there is the risk of including other foreign material (e.g., wax or exuviae) that may contaminate the purity of the sample and make the laboratory analyses and results less affordable.

Collecting beebread samples could be useful to:

- Detect residues of pollutants in the environment such as heavy metals, agrochemicals and varroacides (Conti & Botrè, 2001; Fleche, Clement, Zeggane, & Faucon, 1997; Gooley, Gooley, & Fell, 2018; Kubik et al., 1999; Leita, Muhlbachova, Cesco, Barbattini, & Mondini, 1996; Mullin et al., 2010; Orantes-Bermejo, Pajuelo, Megías, & Fernández-Piñar, 2010; Sattler et al., 2016).
- Evaluate the qualitative and quantitative composition of pollen collected by the bees.
- Verify the presence of toxic or allergenic pollens.

- Detect the presence of bee pathogens (e.g., spores of *Paenibacillus larvae*, *Melissococcus plutonius*, *Ascosphaera apis*, *Nosema* spp.) (Chen, Evans, & Feldlaufer, 2006; Gilliam, Prest, & Lorenz, 1989; Gilliam, Taber, Lorenz, & Prest, 1988; Hale & Menapace, 1980; Higes, Martín-Hernández, Garrido-Bailon, García-Palencia, & Meana, 2008; Mehr, Menapace, Wilson, & Sackett, 1976; Moffet, Wilson, Stoner, & Wardecker, 1978).

A New Tool for Sampling Beebread: the “Beebread Collector”

The purpose of this article is to describe a very simple and low cost tool designed to allow the easy extraction of beebread from comb cells (Figure 2A, B), invented by Giulio Loglio. It is intended to be home-made by interested beekeepers, such as those participating in citizen science projects, but also for those harvesting




Figure 1. (A) Corbicular pollen loads collected using a pollen trap. (B) Beebread cells stored in comb.




Figure 2. (A) The ready assembled beebread collector. (B) Components of the beebread collector: a small rigid tube; a plunger and a spring. (C) How to collect beebread: it is important to insert the beebread collector vertically into the cell. (D) A small pellet of beebread has been expelled from the beebread collector.

beebread for commercial use. The beebread collector can be made of various materials that can be in contact with food avoiding contamination of the beebread (glass, plastic, and steel). The main components of the tool are:

- A rigid tube, several centimeters long, with a diameter between 4.8 and 5.15 mm, the same as the honey bee cells. Commonly available drinking straws often have an external diameter of 4.8 mm. The tube, which has very thin walls, has a flattened top end.
- A plunger, such as a large steel nail with the point cut off, with a diameter of about 4.5 mm, which can slide into the rigid tube. The plunger has a flattened top end and must have a length so as to protrude from the rigid tube for 1–2 mm when fully depressed.
- A small spring 2–3 cm long inserted into the plunger able to bring the plunger in back position when it is not pressed.

To use the tool, the beebread collector is grasped like a pen and inserted vertically into a cell containing the beebread, exerting at the same time both a light pressure and rotary movement to allow the rigid tube to reach the bottom of the cell (Figure 2C). The outer part of the tube, sliding in contact with the inner walls of the cell, detaches the beebread from the cell wall and conveys it into the central part of the tube. To extract the beebread from the collector, simply press the plunger. The beebread is then pushed out of the tube in the form of a small pellet (Figure 2D).

Usually each cell of a bee comb contains about 75 mg of beebread and on average with the beebread collector it is possible to sample about 50 mg of it. To collect 10 g of beebread (the amount that is often required for laboratory analysis), it is thus necessary to insert the beebread collector into about 200–250 cells. The beebread collector can be cleaned with hot water and a small brush (e.g., dental brush), but if necessary to avoid cross contamination, due to its low cost it can be disposed and replaced by a new one.

It is possible to make the beebread collector more efficient and rapid by connecting the rigid tube directly to a vacuum pump so that the beebread, after being extracted from the cell, is deposited in a hermetic container. Two different solutions could be adopted:

1. An electric motor rotates the rigid tube which, inserted in the cell and acting like a small drill or a small cutter, crumbles the beebread.
2. An electric motor rotates in the rigid tube a small metal rod which, ending




Figure 3. Beebread collector equipped with a small electric motor.



Figure 4. Differences between the beebread extracted with the manual beebread collector (pellets, on the left) and the motorized beebread collector (crumbles, on the right).

with a small corkscrew, crumbles the beebread (Figure 3).

In both solutions the beebread, vacuumed with a vacuum pump, is deposited in a hermetic container in a crumbled form and not in the form of pellets (Figure 4).

Conclusions

Beebread is an underused hive product that could have greater value both as a novel food and as a matrix for laboratory analysis to detect residues of agrochemicals, veterinary medicines used by beekeepers, presence of heavy metals in the environment or the presence of several honey bee pathogens. Our beebread collector is a tool that allows sampling in a fast, clean and uniform way, without altering the bee comb structure and guaranteeing the quality of samples for laboratory analysis. This simple and cheap tool will be used in the new INSIGNIA project, which is an innovative monitoring program using beekeepers as citizen scientists to study pesticide exposure of honey bee colonies and investigation of pollen sources.

Acknowledgements


This publication is part of the project “Citizen science investigation for pesticides in apicultural products” (INSIGNIA) – Multi Call PP-1-1-2018 – which has received funding from the European Union. Grant Agreement number: SANTE/E4/SI2.788418 – INSIGNIA – PP-1-1-2018. The content of this publication represents the views of the Authors only and is their responsibility; it cannot be considered to reflect the views of the European Commission or any other body of the European Union. The European Commission does not accept any responsibility for use that may be made of the information it contains.

References

- Andelković, B., Jevtić, G., Mladenović, M., Marković, J., Petrović, M., & Nedić, N. (2012). Quality of pollen and honey bee bread collected in spring. *Journal of Hygienic Engineering and Design*, 1, 275–277.
- Anderson, K. E., Carroll, M. J., Sheehan, T. I. M., Mott, B. M., Maes, P., & Corby-Harris, V. (2014). Hive-stored pollen of honey bees: Many lines of evidence are consistent with pollen preservation, not nutrient conversion. *Molecular Ecology*, 23(23), 5904–5917. doi:10.1111/mec.12966
- Barajas, J., Cortes-Rodriguez, M., & Rodríguez-Sandoval, E. (2012). Effect of temperature on the drying process of bee pollen from two zones of Colombia. *Journal of Food Process Engineering*, 35(1), 134–148. doi:10.1111/j.1745-4530.2010.00577.x
- Barene, I., Daberte, I., & Silksna, S. (2015). Investigation of bee bread development of its dosage forms. *Medicinas teorija it practika*, 21(1), 16–22. doi:10.15591/mtp.2015.003
- Campos, M. G., Bogdanov, S., de Almeida-Muradian, L. B., Szczesna, T., Mancebo, Y., Frigerio, C., & Ferreira, F. (2008). Pollen composition and standardisation of analytical methods. *Journal of Apicultural Research*, 47(2), 154–161. doi:10.1080/00218839.2008.11101443
- Carroll, M. J., Brown, N., Goodall, C., Downs, A. M., Sheehan, T. H., & Anderson, K. E. (2017). Honey bees preferentially consume freshly-stored pollen. *PLoS One*, 12(4), e0175933.
- Chen, Y., Evans, J., & Feldlaufer, M. (2006). Horizontal and vertical transmission of viruses in the honey bee, *Apis mellifera*. *Journal of Invertebrate Pathology*, 92, 152–159. doi:10.1016/j.jip.2006.03.010
- Conti, M. E., & Botrè, F. (2001). Honeybees and their products as potential bioindicators of heavy metals contamination. *Environmental Monitoring and Assessment*, 69(3), 267–282. doi:10.1023/A:1010719107006
- De Grandi-Hoffman, G., Chen, Y., & Simonds, R. (2013). The effects of pesticides on queen rearing and virus titers in honey bees (*Apis mellifera* L.). *Insects*, 4(1), 71–89.
- Estevinho, M. L., Afonso, S. E., & Feás, X. (2011). Antifungal effect of lavender honey against *Candida albicans*, *Candida krusei* and *Cryptococcus neoformans*. *Journal of Food Science and Technology*, 48(5), 640–643. doi:10.1007/s13197-011-0243-1
- Fleche, C., Clement, M. C., Zeggane, S., & Faucon, J. P. (1997). Contamination of bee products and risk for human health: Situation in France. *Revue scientifique et technique (International Office of Epizootics)*, 16(2), 609–619. doi:10.20506/rst.16.2.1049
- Fuenmayor, B., Zuluaga, D., Díaz, M., Quicazán de C, M., Cosio, M., & Mannino, S. (2014). Evaluation of the physicochemical and functional properties of Colombian bee pollen. *Revista MVZ Córdoba*, 19(1), 4003–4014. doi:10.21897/rmvz.120
- Gilliam, M. (1979). Microbiology of pollen and bee bread: The yeasts. *Apidologie*, 10(1), 43–53. doi:10.1051/apido:19790106
- Gilliam, M., Prest, D. B., & Lorenz, B. J. (1989). Microbiology of pollen and bee bread: Taxonomy and enzymology of molds. *Apidologie*, 20, 53–68. doi:10.1051/apido:19890106
- Gilliam, M., Taber III, S., Lorenz, B. J., & Prest, D. B. (1988). Factors affecting development of chalkbrood disease in colonies of honey bees, *Apis mellifera*, fed pollen contaminated with *Ascosphaera apis*. *Journal of Invertebrate Pathology*, 52, 314–325. doi:10.1016/0022-2011(88)90141-3
- Gooley, Z. C., Gooley, A. C., & Fell, R. D. (2018). Relationship of landscape type on neonicotinoid insecticide exposure risks to honey bee colonies: A statewide survey. *Journal of Economic Entomology*, 111(6), 2505–2512. doi:10.1093/jeet/toy284.
- Grodzinski, W., & Yorks, T. P. (1981). Species and ecosystem bioindicators of airborne pollution: An analysis of two major studies. *Water, Air and Soil Pollution*, 16, 33–53. doi:10.1007/BF01047040
- Habryka, C., Kruczek, M., & Drygaś, B. (2016). Bee products used in apitherapy. *World Scientific News*, 48, 254–258.
- Hale, P. J., & Menapace, D. M. (1980). Effect of time and temperature on the viability of *Ascosphaera apis*. *Journal of Invertebrate Pathology*, 36, 429–430. doi:10.1016/0022-2011(80)90050-6
- Higes, M., Martín-Hernández, R., Garrido-Bailon, E., García-Palencia, P., & Meana, A. (2008). Detection of infective *Nosema ceranae* (Microsporidia) spores in corbicular pollen of forager honeybees. *Journal of Invertebrate Pathology*, 97(1), 76–78. doi:10.1016/j.jip.2007.06.002
- Karmakar, P. (2015). Pollen analysis in understanding the foraging behaviour of bee species used for apicultural practice in gangetic West Bengal.
- Kieliszek, M., Piowarek, K., Kot, A. M., Błażej, S., Chlebowska-Śmigiel, A., & Wolska, I. (2017). Pollen and bee bread as new health-oriented products: A review. *Trends in Food Science & Technology*, 71, 170–180. doi:10.1016/j.tifs.2017.10.021
- Kubik, M., Nowacki, J., Pidek, A., Warakomska, Z., Michalczuk, L., & Goszczynski, W. (1999). Pesticide residues in bee products collected from cherry trees protected during blooming period with contact and systemic fungicides. *Apidologie*, 30(6), 521–532. doi:10.1051/apido:19990607
- Leita, L., Muhlbachova, G., Cesco, S., Barbattini, R., & Mondini, C. (1996). Investigation of the use of honey bees and honey bee products to assess heavy metals contamination. *Environmental Monitoring and Assessment*, 43(1), 1–9. doi:10.1007/BF00399566
- Mehr, Z., Menapace, D. M., Wilson, W. T., & Sackett, R. R. (1976). Studies on the initiation and spread of chalkbrood within an apiary. *American Bee Journal*, 116, 266–268.
- Moffet, J. O., Wilson, W. T., Stoner, A., & Wardecker, A. (1978). Feeding commercially purchased pollen containing mummies caused chalkbrood. *American Bee Journal*, 118, 412–414.
- Mullin, C. A., Frazier, M., Frazier, J. L., Ashcraft, S., Simonds, R., & Pettis, J. S. (2010). High levels of miticides and agrochemicals in North American apiaries: Implications for honey bee health. *PLoS One*, 5(3), e9754.
- Nagai, T., Nagashima, T., Myoda, T., & Inoue, R. (2004). Preparation and functional properties of extracts from bee bread. *Food/Nahrung*, 48(3), 226–229.
- Orantes-Bermejo, F. J., Pajuelo, A. G., Megías, M. M., & Fernández-Piñar, C. T. (2010). Pesticide residues in beeswax and beebread samples collected from honey bee colonies (*Apis mellifera* L.) in Spain. Possible implications for bee losses. *Journal of Apicultural Research*, 49(3), 243–250. doi:10.3896/IBRA.1.49.3.03
- Porrini, C., Ghini, S., & Girotti, S. (2002). Use of honey bees as bioindicators of environmental pollution in Italy. In Devillers J., Pham-Delègue M.H. (Eds.), *Honey bees: Estimating the environmental impact of chemicals* (pp. 186–247). London and New York: Taylor & Francis.
- Raes, H., Cornelis, R., & Rzezniak, U. (1992). Distribution, accumulation and depuration of administered lead in adult honey bees. *Science of the Total Environment*, 113, 269–279. doi:10.1016/0048-9697(92)90005-D
- Sattler, J. A. G., De-Melo, A. A. M., Nascimento, K. S. D., Mancini-Filho, J., Sattler, A., & Almeida-Muradian, L. B. D. (2016). Essential minerals and inorganic contaminants (barium, cadmium, lithium, lead and vanadium) in dried bee pollen produced in Rio Grande do Sul State, Brazil. *Food Science and Technology*, 36(3), 505–509.
- van der Steen, J. J. M., Cornelissen, B., Blacquièrre, T., Pijnenburg, J. E. M. L., & Severijnen, M. (2016). Think regionally, act locally: Metals in honeybee workers in the Netherlands (surveillance study 2008). *Environmental Monitoring and Assessment*, 188(8), 463.
- van der Steen, J. J. M. (2016). *Beehold: The colony of the honeybee (Apis mellifera L.) as a bio-sampler for pollutants and plant pathogens* (Doctoral dissertation). Wageningen University.
- Vásquez, A., & Olofsson, T. C. (2009). The lactic acid bacteria involved in the production of bee pollen and bee bread. *Journal of Apicultural Research*, 48(3), 189–195. doi:10.3896/IBRA.1.48.3.07
- Zuluaga, C. M., Serrato, J. M., & Quicazan, M. C. (2015). Chemical, nutritional and bioactive characterization of colombian bee-bread. *Chemical Engineering Transactions*, 43, 175–180.

Giulio Loglio
Veterinarian and Beekeeper, Italy

Giovanni Formato
Istituto Zooprofilattico Sperimentale del Lazio e della Toscana “M. Aleandri”, Italy
Email: giovanni.formato@izslt.it
 <http://orcid.org/0000-0002-1202-5745>

Marco Pietropaoli
Istituto Zooprofilattico Sperimentale del Lazio e della Toscana “M. Aleandri”, Italy
 <http://orcid.org/0000-0002-9073-909X>

Riccardo Jannoni-Sebastianini
International Federation of Beekeepers’ Association, APIMONDIA, Italy

Norman Carreck
Carreck Consultancy Ltd., UK

Jozef van der Steen
Alveus AB Consultancy, the Netherlands