



UNIVERSITÀ
DI SIENA
1240



6TH APIMEDICA

&

22-25 NOVEMBER 2016 ROME

5TH APIQUALITY

**Bee propolis in the
treatment of
Helicobacter pylori:
in the right way to
clinical application**


Marco Biagi

**UNIVERSITÀ DEGLI STUDI DI SIENA
SOCIETÀ ITALIANA DI FITOTERAPIA**




Helicobacter pylori

Helicobacter pylori infection is the most widespread infection and over a billion of people worldwide are affected.



Helicobacter pylori




Isolated in 1982, *Helicobacter pylori* is a Gram- flagellated bacterium, able to live in the human stomach acidic and microaerophilic environment.

HP have **hydrogenases** that convert molecular hydrogen of intestinal flora in energy, **catalases**, **oxidases** and **ureases**.

Urease convert urea in carbon dioxide and ammonia increasing pH and facilitating the colonization of the bacterium.




Helicobacter pylori: clinical evolution




Helicobacter pylori: pathogenesis

The most important gastric diseases related to HP infections are:

- Gastritis
- Peptic ulcer
- Metaplasia
- Dyspepsia
- Gastric carcinoma
- MALT Lymphoma



Helicobacter pylori: pathogenesis



Helicobacter pylori: virulence factors

Pathogenicity:

- **Adhesins (BabA/B, SabA)**: penetration and adhesion to the gastric mucosa
- **Bacterial lipopolysaccharide**: massive inflammatory response
- **Urease**: adhesion factor, nitrogen source, human DNA damage, increase of inflammatory response
- **HP-NAP** (HP neutrophil activating protein): *increase of ROS and cytokines release.*
- **VacA protein**: *vacuolizant and cytotoxic protein. VacA alters cell tight junctions, increase Ni^{2+} , Fe^{3+} ions and other bacterial nutrient influx.*
- **CagA protein**: *massive inflammatory response (IL-6, IL-8, IL-18, IL-1 β release) and macrophages activation. Alteration of gastric epithelial cells proliferation.*

Helicobacter pylori: conventional treatment

According to **Maastricht/Florence V Consensus 2016** guidelines, the conventional anti-HP treatment is:

I Line: 14 days

PPI
Clarithromycin 500 mg x 2
Amoxicillin 1 g x 2

or

PPI
Clarithromycin 500 mg x 2
Metronidazole 500 mg x 2

Failure

II Linea: 14 days

PPI
Bismuth salts 120 mg
Metronidazole 500 mg
Tetracycline 500 mg

o

PPI
Tetracycline 500 mg x 2
Metronidazole 500 mg x 2

o

PPI
Amoxicillin 1 g x 2
Metronidazole 500 mg x 2

If failure occur: Levofloxacin- containing triple or quadruple therapy may be recommended

Helicobacter pylori treatment: issues and limits



- ***Eradication rate not > 80-85%***
- Metronidazole resistance (>40%)
- Clarithromycin (>20%)
- **Side effects (gut flora damage, fatigue)**



Helicobacter pylori treatment: issues and limits

HP eradication do not resolve inflammatory symptoms in the first weeks after treatment.

- > 50% of subjects with gastric ulcer maintain symptoms;
- Oxidative stress.



Propolis in the Helicobacter pylori treatment

Rationale of the choice:

- Clinical investigations refer the effectiveness of propolis against *Helicobacter pylori* infections.
- Antioxidant and anti-inflammatory properties
- Safety profile



Our work


1-Comparison of different propolis marketed in Italy.

2-Evaluation of the effectiveness against several HP strains.

3-gastric stability.

4-Evaluation of the synergy between propolis and anti-HP antibiotics.

5-*in vitro* anti-inflammatory activity.



Tested samples

Propolis dried extract (12-15% total flavonoids) (**WIPE**)

Hydrodispersible propolis dried extract (8-10% total flavonoids) (**HDPE**)

Standardized 50% total polyphenols propolis based dried extract (**FEPE**)



Chemical analyses and antiradicalic and antioxidant *in vitro* essays

sample	Total polyphenols %	Flavonoids %	IC50 (DPPH) μg/ml	Ox. Pot. V
FEPE	50,82% ± 1,66	11,23% ± 0,84%	10,2 ± 0,9	0,30 – 0,50
HDPE	18,08% ± 2,64	11,47% ± 0,11%	10,4 ± 0,6	0,30 – 0,57
WIPE	18,95% ± 3,36	12,02% ± 0,67%	65,4 ± 1,4	0,32 – 0,52



anti *Helicobacter pylori* activity

<i>Prop.</i>	<i>MIC mg/l</i>	<i>MBC mg/l</i>	<i>HP strain</i>
HDPE	>2500	>2500	10K VacA+ CagA+
FEPE	625	1250	
WIPE	1250	2500	
HDPE	>2500	>2500	G21 VacA+ CagA-
FEPE	625	1250	
WIPE	625	1250	
HDPE	>2500	>2500	G27 clar. resistant
FEPE	625	1250	
WIPE	1250	2500	
HDPE	>2500	>2500	328 IC gastric carcinoma
FEPE	625	1250	
WIPE	625	1250	
HDPE	>2500	>2500	CCUG metron. resistant
FEPE	625	1250	
WIPE	1250	2500	

MIC: minimum inhibitory concentration

MBC: minimum bactericidal concentration



Influence of bacterial concentration on FEPE propolis extract activity

Strain :
CCUG metron. Res.

Sample:
FEPE

CFU/ml	MIC mg/l	MBC mg/l
10^3	625	1250
10^6	625	1250
10^9	625	1250

No MIC or MBC variations .



Killing Time

Strain :
CCUG metron. Res.

Sample:
FEPE

Time	MBC mg/l
2 h	--
4 h	1250
24 h	1250

Propolis is effective against HP in 4 h.

The mechanism of action is probably related to a physical bacterial wall damage.

≠ biochemical mechanism of antibiotics



Gastric stability of propolis extract

According to USP e McDougall 2006 (modified)

pH	T ₀ polyphenols mg/l	post digestion polyphenols mg/l	recovery
1,7	418,48 ± 2,81	384,76 ± 24,95	91,20% ± 5,86%
4,5	418,48 ± 2,81	377,93 ± 25,50	89,36% ± 5,99%



Antibacterial synergy between propolis and antibiotics

Antibiotics:

Clarithromycin, 1 mg/l

Metronidazole, 64 mg/l

Tetracycline, 8 mg/l

Levofloxacin, 16 mg/l

Amoxicillin, 2 mg/l

Strains:

1-CCUG, metronidazole resistant

2- G27, clarithromycin resistant

Propolis extract:

FEPE




Checkerboard method

200 μ L FEPE in BB



100 μ L
antib.



*Evaluation of
FIC and FBC indexes
according to EUCAST*



Antibacterial synergy between propolis and antibiotics

Propolis and Clarithromycin

CCUG metron.res.	FIC	FBC	FIC index	FBC index
Propolis	n.d	n.d	n.d	n.d
Clarithromycin	<0,250	<0,250		
IC clarit. res.	FIC	FBC	FIC index	FBC index
Propolis	0,250	0,250	0,500	0,500
Clarithromycin	0,250	0,250		



Antibacterial synergy between propolis and antibiotics

Propolis and Metronidazole

CCUG metron.res.	FIC	FBC	FIC index	FBC index
Propolis	0,250	0,500	0,500	0,531
Metronidazole	0,250	0,031		
IC clarit. res.	FIC	FBC	FIC index	FBC index
Propolis	0,500	0,500	0,750	0,750
Metronidazole	0,250	0,250		



Antibacterial synergy between propolis and antibiotics

Propolis and Tetracycline

CCUG metron.res.	FIC	FBC	FIC index	FBC index
Propolis	0,500	1,000	0,625	1,031
Tetracycline	0,125	0,031		
IC clarit. res.	FIC	FBC	FIC index	FBC index
Propolis	0,500	0,500	0,516	0,516
Tetracycline	0,016	0,016		



Antibacterial synergy between propolis and antibiotics

Propolis and Levofloxacin

CCUG metron.res.	FIC	FBC	FIC index	FBC index
Propolis	0,250	0,125	0,375	0,188
Levofloxacin	0,125	0,063		
IC clarit. res.	FIC	FBC	FIC index	FBC index
Propolis	1,000	1,000	1,500	1,500
Levofloxacin	0,500	0,500		



Antibacterial synergy between propolis and antibiotics

Propolis and Amoxicillin

CCUG metron.res.	FIC	FBC	FIC index	FBC index
Propolis	0,250	0,250	<0,266	0,258
Amoxicillin	≤0,016	0,008		
IC clarit. res.	FIC	FBC	FIC index	FBC index
Propolis	0,250	0,250	0,500	0,500
Amoxicillin	0,250	0,250		



Antibacterial synergy between propolis and antibiotics

- Bacteriostatic synergy** (FIC indexes) occur in **60% of tested situations**;
- Bactericidal synergy** (FBC indexes) occur in **50% of cases**.


In 30% of cases FIC_i and FBC_i show that propolis and antibiotics effects are additive.

In worst cases propolis and antibiotics have indifferent synergy.



ex vivo PBMC anti-inflammatory activity

Modulazione dell'espressione della COX-2



Discussion

Our findings showed that propolis could be used conveniently in clinical practices in HP infections management, in particular in association with conventional PPI-antibiotic therapy.

1- HP loss of resistance

2- Synergistic effect of prpolis and antibiotics

3- Antioxidant and anti-inflammatory effectiveness

Propolis integration in the HP treatment could increase eradication rate, help in HP related symptoms, decrease side effects and have positive ripercussions on patients' compliance.

